

Subpart B—Specific Tolerances for Residues of New Animal Drugs

§ 556.20 2-Acetylamino-5-nitrothiazole.

A tolerance of 0.1 part per million is established for negligible residues of 2-acetylamino-5-nitrothiazole in the edible tissues of turkeys.

§ 556.30 Aklomide.

Tolerances are established for combined residues of aklomide (2-chloro-4-nitrobenzamide) and its metabolite (4-amino-2-chlorobenzamide) in uncooked edible tissues of chickens as follows:

- (a) 4.5 parts per million in liver and muscle.
- (b) 3 parts per million in skin with fat.

§ 556.34 Albendazole.

Tolerances are established for residues of albendazole in uncooked edible tissues as follows:

(a) *Cattle*. The tolerance for the 2-aminosulfone metabolite (marker residue) in cattle liver (target tissue) is 0.2 part per million. The tolerance refers to the concentration of marker residue in the target tissue used to monitor for total drug residues in the target animals.

(b) *Sheep*. The tolerance for the 2-aminosulfone metabolite (marker residue) in sheep liver (target tissue) is 0.25 part per million.

[59 FR 65711, Dec. 21, 1994]

§ 556.38 Amoxicillin.

A tolerance of 0.01 part per million is established for negligible residues of amoxicillin in milk and in the uncooked edible tissues of cattle.

[49 FR 45422, Nov. 16, 1984]

§ 556.40 Ampicillin.

A tolerance of 0.01 p/m is established for negligible residues of ampicillin in the uncooked edible tissues of swine and cattle and in milk.

§ 556.50 Amprolium.

Tolerances are established as follows for residues of amprolium (1-(4-amino-2-*n*-propyl-5-pyrimidinylmethyl)-2-picolinium chloride hydrochloride):

(a) In the edible tissues and in eggs of chickens and turkeys:

(1) 1 part per million in uncooked liver and kidney.

(2) 0.5 part per million in uncooked muscle tissue.

(3) In eggs:

(i) 8 parts per million in egg yolks.

(ii) 4 parts per million in whole eggs.

(b) In the edible tissues of calves:

(1) 2.0 parts per million in uncooked fat.

(2) 0.5 part per million in uncooked muscle tissue, liver, and kidney.

(c) In the edible tissues of pheasants:

(1) 1 part per million in uncooked liver.

(2) 0.5 part per million in uncooked muscle.

[40 FR 13942, Mar. 27, 1975, as amended at 50 FR 18472, May 1, 1985]

§ 556.52 Apramycin.

A tolerance of 0.1 part per million is established for parent apramycin (marker residue) in kidney (target tissue) of swine. The acceptable daily intake (ADI) for total residues of apramycin is 25 micrograms per kilogram of body weight per day.

[62 FR 40933, July 31, 1997]

§ 556.60 Arsenic.

Tolerances for total residues of combined arsenic (calculated as As) in food are established as follows:

(a) In edible tissues and in eggs of chickens and turkeys:

(1) 0.5 part per million in uncooked muscle tissue.

(2) 2 parts per million in uncooked edible by-products.

(3) 0.5 part per million in eggs.

(b) In edible tissues of swine:

(1) 2 parts per million in uncooked liver and kidney.

(2) 0.5 part per million in uncooked muscle tissue and by-products other than liver and kidney.

§ 556.70 Bacitracin.

Tolerances for residues of bacitracin from zinc bacitracin or bacitracin methylene disalicylate are established at 0.5 part per million (0.02 unit per gram), negligible residue, in uncooked

edible tissues of cattle, swine, chickens, turkeys, pheasants, and quail, and in milk and eggs.

[42 FR 18614, Apr. 8, 1977]

§ 556.90 Buquinolate.

Tolerances are established for residues of buquinolate as follows:

- (a) In edible tissues of chickens:
 - (1) 0.4 part per million in uncooked liver, kidney, and skin with fat.
 - (2) 0.1 part per million in uncooked muscle.
- (b) In eggs:
 - (1) 0.5 part per million in uncooked yolk.
 - (2) 0.2 part per million in uncooked whole eggs.

§ 556.100 Carbadox.

A tolerance of 30 parts per billion is established for residues of quinoxaline-2-carboxylic acid (marker residue) in liver (target tissue) of swine.

[63 FR 13337, Mar. 19, 1998]

§ 556.110 Carbomycin.

A tolerance of zero is established for residues of carbomycin in the uncooked edible tissues of chickens.

§ 556.113 Ceftiofur.

Cattle, swine, poultry, and sheep: A tolerance for residues of ceftiofur in edible tissue is not required.

[57 FR 41862, Sept. 14, 1992, as amended at 61 FR 66583, Dec. 18, 1996]

§ 556.115 Cephapirin.

A tolerance of 0.02 parts per million (ppm) is established for residues of cephalapirin in the milk and 0.1 ppm in the uncooked edible tissues of dairy cattle.

[40 FR 57454, Dec. 10, 1975]

§ 556.120 Chlorhexidine.

A tolerance of zero is established for residues of chlorhexidine in the uncooked edible tissues of calves.

§ 556.140 Chlorobutanol.

A tolerance of zero is established for residues of chlorobutanol in milk from dairy animals.

§ 556.150 Chlortetracycline.

Tolerances are established for the sum of residues of the tetracyclines including chlortetracycline, oxytetracycline, and tetracycline, in tissues of beef cattle, nonlactating dairy cows, calves, swine, sheep, chickens, turkeys, and ducks, as follows:

- (a) 2 parts per million (ppm) in muscle.
- (b) 6 ppm in liver.
- (c) 12 ppm in fat and kidney.

[61 FR 67453, Dec. 23, 1996]

§ 556.160 Clopidol.

Tolerances for residues of clopidol (3,5-dichloro-2,6-dimethyl-4-pyridinol) in food are established as follows:

- (a) In cereal grains, vegetables, and fruits: 0.2 part per million.
- (b) In chickens and turkeys:
 - (1) 15 parts per million in uncooked liver and kidney.
 - (2) 5 parts per million in uncooked muscle.
- (c) In cattle, sheep, and goats:
 - (1) 3 parts per million in uncooked kidney.
 - (2) 1.5 parts per million in uncooked liver.
 - (3) 0.2 part per million in uncooked muscle.
- (d) In swine: 0.2 part per million in uncooked edible tissues.
- (e) In milk: 0.02 part per million (negligible residue).

§ 556.163 Clorsulon.

Tolerances are established for residues of clorsulon in cattle as follows:

- (a) The tolerance for clorsulon (market residue) in kidney (target tissue) is 1.0 part per million. A marker residue of 1.0 part per million corresponds to a total residue of 3.0 parts per million in kidney.
- (b) The safe concentrations for total clorsulon residues in uncooked edible cattle tissues are: muscle, 1.0 part per million; liver, 2.0 parts per million; kidney, 3.0 parts per million; and fat, 4.0 parts per million.

[50 FR 10221, Mar. 14, 1985]

§ 556.165 Cloxacillin.

A tolerance of 0.01 part per million is established for negligible residues of